

# DISINFECTION OF FIELD EQUIPMENT TO MINIMIZE RISK OF SPREAD OF CHYTRIDIOMYCOSIS AND RANAVIRUS<sup>1</sup>

## IMPORTANCE OF DISINFECTION

The spread of pathogens is a major threat to amphibians and reptiles worldwide.<sup>2-5</sup> This is particularly true for Ranavirus (RV) and *Batrachochytrium dendrobatidis* (Bd) responsible for chytridiomycosis. Humans can transmit diseases from one place to another and from one organism to another in a short amount of time and over distances the organisms cannot traverse. With the increasing spread of pathogens and reports of die-offs among amphibians and select reptiles worldwide, it is imperative that field biologists, researchers, hobbyists, and anyone interested in recreational herpetology-related field activities employ basic disinfecting procedures to prevent the spread of pathogens.

## BEFORE LEAVING FOR THE FIELD

Although other chemicals are effective (see table), NEPARC recommends a 3% bleach solution to inactivate Bd and most RV's.<sup>3-7</sup> Concentrated bleach is inexpensive and readily available. However, diluted bleach solutions lose their potency if exposed to air, sunlight, or organic material, and should be discarded after 5 days if exposed.<sup>8</sup> To ensure maximum efficacy, prepare only as much solution as you will need for the sampling event.

### Suggested equipment:

- Brushes for scrubbing and/or removing mud and vegetation from equipment.
- Hand sanitizers and antiseptic alcohol wipes.
- Handheld bottles and/or pump sprayers for applying bleach and water. Bring clean rinse water.



- Gloves for handling animals. These should be disinfected or discarded between animals.
- Plastic bags of different sizes: examining animals in bag minimizes contact.
- Prepare additional sets of equipment if sampling at multiple locations.
- Trash bags.

## INSTRUCTIONS FOR LARGE EQUIPMENT

Brush off mud, wash with biodegradable soap, disinfect with bleach and rinse all exterior surfaces of boats, canoes, vehicles or trailers and their tires that may have come in contact with potentially affected water (e.g. stream or wetland).

## AFTER EACH SAMPLING EVENT AND BEFORE MOVING TO THE NEXT SITE

1. Brush off mud and vegetation from field equipment (e.g., nets, buckets, boots). Soil or mud can reduce the effectiveness of the disinfection process.
2. Generously spray or immerse all items in bleach solution.
  - Bleach is highly toxic to aquatic organisms; stand at least 50 m from any natural water source.
  - Lab studies indicate 1 minute contact time to be sufficient to inactivate pathogens but NEPARC recommends 5 minutes in field situations.
3. Rinse bleached items with water to minimize damage to the equipment and to prevent exposing the next wetland to residual bleach.
4. Use alcohol wipes to disinfect calipers, measuring boards, and other sensitive equipment.



## END OF THE DAY

After returning from the field, all equipment should be washed and thoroughly disinfected. If available, set up 2 buckets or large tubs: one with soapy water and one with 3% bleach solution.

- Brush or scrub off any soil or vegetation. Immerse into soap, wash then rinse.
- Immerse in bleach and leave for 5 minutes. Rinse thoroughly with water.
  - Hang equipment and gear, and allow them to air dry completely.

# DISINFECTION OPTIONS FOR RANAVIRUS (RV) AND *BATRACHOCHYTRIUM DENDROBATIDIS* (Bd)

Although these chemicals were not developed specifically for RV or Bd, these recommendations represent the minimum concentration and contact time demonstrated as effective

	<b>Clorox Bleach®</b>	<b>Nolvasan®</b>	<b>Virkon S®</b>	<b>Ethanol</b>
<b>Active Ingredient (AI)</b>	Sodium hypochlorite	Chlorhexidine	Potassium peroxydisulfate	Ethyl alcohol
<b>Concentration of AI</b>	6.0%	2.0%	20.4%	70.0%
<b>Relative cost</b>	\$4.99/gal	\$65.95/gal	\$76.50/10 lb or \$1.60/gal	\$23.45/L or \$88.83/gal
<b>Min. Contact Time RV<sup>9</sup>/Bd<sup>10</sup></b>	1 min / 30 sec	1 min / not determined	1 min / 20 sec	1 min <sup>11</sup> / 20 sec
<b>Min. Concentration RV<sup>9</sup>/Bd<sup>10</sup></b>	3.0% / 1.0%	0.75% / not determined	1.0% / 1.0%	70% / 70%
<b>Effective dilution ratio for both RV and Bd</b>	1:32 dilution (bleach:water) for 3% solution using 6% concentration of household bleach.	1:127 (Nolvasan®: water) for 0.75% solution (RV only)	1 scoop (1.3 oz) or 1 tablet per gal of water	Effective when applied undiluted (70%)
<b>Toxicity to Humans</b>	<ul style="list-style-type: none"> <li>Vapor may cause severe irritation or damage to eyes and skin</li> <li>Harmful if swallowed</li> </ul>	<ul style="list-style-type: none"> <li>May be fatal if inhaled</li> <li>Avoid breathing spray mist</li> <li>Causes irreversible eye damage</li> <li>Harmful if swallowed</li> </ul>	<ul style="list-style-type: none"> <li>Harmful if swallowed</li> <li>Irritating to respiratory system and skin</li> <li>May cause serious eye damage</li> </ul>	<ul style="list-style-type: none"> <li>May be fatal if swallowed or inhaled</li> <li>Can damage liver, kidneys and nervous system by repeated or prolonged exposure</li> <li>May be absorbed through skin. Repeated or prolonged contact can cause eye irritation or dermatitis<sup>12</sup></li> </ul>
<b>Toxicity to Amphibians</b>	<ul style="list-style-type: none"> <li>Fatal at high concentrations</li> </ul>	<ul style="list-style-type: none"> <li>Safe for short durations<sup>13</sup></li> </ul>	<ul style="list-style-type: none"> <li>Non-toxic<sup>14</sup></li> </ul>	<ul style="list-style-type: none"> <li>May destroy mucus and wax resulting in dehydration and microbial infection<sup>11</sup></li> </ul>
<b>Effects on Equipment</b>	<ul style="list-style-type: none"> <li>Corrodes metals</li> <li>Will fade colors and break down cloth fibers</li> </ul>	<ul style="list-style-type: none"> <li>None reported</li> </ul>	<ul style="list-style-type: none"> <li>Safe on fabric</li> <li>May cause pitting on galvanized or soft metal if not rinsed with water</li> </ul>	<ul style="list-style-type: none"> <li>May damage rubber and plastics</li> <li>May cause deterioration of glues<sup>12</sup></li> </ul>

## Special Instructions:

• Remove debris from equipment prior to treatment.<sup>15</sup> • Wear safety glasses and gloves when handling chemicals. • Water pH can affect chemicals; all information in this table assumes the use of tap or municipal water. • Keep out of lakes, streams, or ponds; stand at least 50 m from any natural water source. • Do not clean equipment or dispose of waste solutions at field sites. • For disposal, follow local, state, and federal guidelines.

**Bleach:** Inactivated by organic material. • Inactivated by sunlight. • If in an opaque container, diluted bleach will last 1 month<sup>16</sup>. If exposed to sunlight or air, it will only last 5 days.

**Nolvasan:** Can be inactivated by organic material.<sup>15</sup> • Store at room temperature in sealed container.<sup>17</sup> • Dilute concentrate with water of pH 5-7.<sup>18</sup> • Remains stable for 1 week if dilute with tap water, and for up to 6 weeks if diluted with deionized water.<sup>17</sup> • Use concentrate within 36 months.<sup>17</sup> • Toxic to fish.<sup>18</sup>

**Virkon-S:** Store at room temperature.<sup>19</sup> • Keep solution away from extreme cold or heat. • Shelf life for tablets is 2 years and for powder is 3 years. • Remains stable for 1 week if diluted with tap water.

**Ethanol:** Highly flammable. • Use and store in a well ventilated area. • Evaporation may diminish effective concentration.<sup>12,18</sup>

## CITATIONS FOR DISINFECTION OF FIELD EQUIPMENT TO MINIMIZE RISK OF SPREAD OF CHYTRIDIOMYCOSIS AND RANAVIRUS

1. This information has been compiled in part from Miller, D. L., and M. J. Gray. 2009. Southeastern Partners in Amphibian and Reptile Conservation, Disease, Pathogens and Parasites Task Team, Information Sheet #10.
2. Converse K.A. and D.E. Green. 2005. Diseases of tadpoles. p 72-88. In: Wildlife Diseases: Landscape Epidemiology, Spatial Distribution and Utilization of Remote Sensing Technology. S.K. Majumdar, J.E. Huffman, F.J. Brenner and A.I. Panah (eds.). Pennsylvania Academy of Science, Easton, PA.
3. Picco, A.M., J.L. Brunner, and J.P. Collins. 2007. Susceptibility of the endangered California tiger salamander, *Ambystoma californiense*, to *Ranavirus* infection. *Journal of Wildlife Diseases* 43:286-290.
4. Picco, A.M. and J.P. Collins. 2008. Amphibian commerce as a likely source of pathogen pollution. *Conservation Biology* 22:1582-1589.
5. St-Amour, V., W.M. Wong, T.W.J. Garner, and D. Lesbarreres. 2008. Anthropogenic influence on prevalence of two amphibian pathogens. *Emerging Infectious Diseases* 14:1175-1176.
6. Bryan L., C.A. Baldwin, M.J. Gray, and D.L. Miller. 2009. Efficacy of select disinfectants at inactivating *Ranavirus*. *Diseases of Aquatic Organisms* 84:89-94.
- 7a. Johnson, M. and R. Speare. 2003. Survival of *Batrachochytrium dendrobatidis* in water: quarantine and control implications. *Emerging Infectious Diseases* 9:922-925.
- 7b. Brem F., J.R. Mendelson III, and K.R. Lips. 2007. Field-Sampling Protocol for *Batrachochytrium dendrobatidis* from Living Amphibians, Using Alcohol Preserved Swabs. Version 1.0. <http://www.amphibianark.org/pdf/>, Accessed 19 March 2014.
8. Green, D.E., M.J. Gray and D.L. Miller. 2009. Disease monitoring and biosecurity. p 481-506. In: *Amphibian Ecology and Conservation: A Handbook of Techniques*. C.K. Dodd (ed.). Oxford University Press, Oxford, United Kingdom.
9. Bryan, L. K., Baldwin, C. A., Gray, M. J., & Miller, D. L. (2009). Efficacy of select disinfectants at inactivating *Ranavirus*. *Diseases of Aquatic Organisms* 84:89-94.
10. Johnson. M., L. Berger, L. Philips, and R. Speare. 2003. Fungicidal effects of chemical disinfectants, UV light, dessication. and heat on the amphibian chytrid, *Batrachochytrium dendrobatidis*. *Diseases of Aquatic Organisms* 57:255-260.
11. Phillott A.D., R. Spear, H.B. Hines, E. Meyer, L.F. Skerratt, K.R. McDonald, S.D. Cashins, D.Mendez, and L. Berger. 2010. Minimising exposure of amphibians to pathogens during field studies. *Diseases of Aquatic Organisms* 92:175-185.
12. Simmons, B., M. Trusler, J. Roccaforte, P. Smith, and R. Scott. 1990. Infection control for home health. *Infection Control and Hospital Epidemiology* 11:362-70.
13. Hadfield, C.A. and B.R. Whitaker. 2005. Amphibian emergency medicine and care. *Seminars in Avian and Exotic Pet Medicine* 14:79-89.
14. Schmidt, B. R., C. Geiser, N. Peyer, N. Keller, and M. von Rutte. 2009. Assessing whether disinfectants against the fungus *Batrachochytrium dendrobatidis* have negative effects on tadpoles and zooplankton. *Amphibia-Reptilia* 30: 313-319.
15. Kennedy, J., J. Bek, and D. Griffin. 2000. Selection and Use of Disinfectants. University of Nebraska Cooperative Extension, G00-1410-A.
16. Rutala, W.A. and J.W. Weber. 1997. Uses of inorganic hypochlorite (bleach) in health care facilities. *Clinical Microbiology Reviews* 10:597-610.

## CITATIONS FOR DISINFECTION OF FIELD EQUIPMENT TO MINIMIZE RISK OF SPREAD OF CHYTRIDIOMYCOSIS AND RANAVIRUS (CONTINUED)

17. Zoetis Veterinary Medical Information and Product Support, inquiry 2014-US-03903, 18 March, 2014.
18. Dvorak, G. 2008. Disinfection 101. Center for Food Security and Public Health. <http://www.cfsph.iastate.edu/Disinfection/Assets/Disinfection101.pdf>, Accessed 19 March 2014.
19. Pharmacal Research Laboratories Inc. <http://www.pharmacal.com/MSDS/US/MSDSVIRKON-S%20Tablet.pdf>, Accessed 19 March 2014.